

## Veterinary Health Requirements for Importation of Cattle

Exporting Country: **Australia**

Importing Country: **Sri Lanka**

**The Government Veterinary Authority of Australia should certify that,**

1. **Australia** is free from Foot and Mouth Disease (FMD) (all serotypes), Vesicular Stomatitis (VS), Contagious Bovine Pleuro- Pneumonia (CBPP), Lumpy Skin Disease(LSD), Rift Valley Fever(RVF), Rabies, Screw worm infestation, Malignant Catarrhal Fever, Theileriosis (*T.Parva*), and Bovine spongiform Encephalopathy (BSE), Haemorrhagic septicaemia (HS). Trypanosomiasis and vaccination against the afore mentioned diseases is not practiced in the country.

### 2. Herd of Origin

- 2.1 The herd of origin of the animals should be free of Bovine Viral Diarrhoea (BVD/MD), Infectious Bovine Rhinotracheitis (IBR/ IPV) and Bovine Tuberculosis, for three (3) years and Johne's disease (Partuberculosis) for six (6) years immediately prior to embarkation. Herd should be non-vaccinated for said diseases.
- 2.2. The herd of origin of the animals should be free of Mycoplasma bovis, Chlamydiosis, Anthrax, Bluetongue, Q Fever, Epizootic haemorrhagic disease(EHD) Salmonellosis caused by *Salmonella Dublin*, Enzootic Bovine Leukosis (EBL), Bovine Genital Campylobacteriosis, Trichomoniasis, Brucellosis (*B.abortus*, *B.melitensis*, *B.Suis*) Leptospirosis and Dermatophilosis, for three (3) years and immediately prior to and till the scheduled date of export.
- 2.3. The herd of origin of the animals should be free of Bovine Babesiosis (*B.bovis*, *B.bigemina*), Bovine Anaplasmosis (*A.marginale* and *A.centrale*), for the last twelve (12) months immediately prior to embarkation.

### 3. Détails of the Animal

- 3.1. The animals intended for export to be kept since birth in Australia.
- 3.2. The selected animals must be AI born with semen originating from government approved semen collection centres.
- 3.3. The animals intended for export are individually identified with a Radio Frequency Identification Device (RFID) approved in Australia under the National Identification and Tracing scheme (NAIT) and under the conditions laid down by OIE (chapter 4.3 of Terrestrial Animal Health Code 2019) to trace back for the information on health, production and genetic background related to the origin of the animals.
- 3.4. The animals have never been fed ruminant protein derived from BSE reported countries except milk protein.
- 3.5. The animals to be non- pregnant, non- vaccinated heifers and tested within 21 days immediately prior to shipment and found to be negative for both antibody and antigen of BVD and IBR.

- 3.6. The animals have not been in contact with any animal showing clinical signs of any infectious or contagious diseases for a period of three months immediately prior to embarkation.

#### 4. Testing and vaccination of Animals

- 4.1. During the pre-export quarantine period the individual animals to be tested by caudal fold test or comparative cervical test for **Bovine tuberculosis**, with negative results.
- 4.2. The individual animals to be tested for both antibody and antigen and found to be negative for **BVD/MD** and **IBR/IPV** during the pre-export quarantine period using ELISA not less than twenty one (21) days and not more than three (3) months prior to embarkation
- 4.3. Animals to be tested for antibody of **Blue tongue** by ELISA and found to be negative during pre-export quarantine period.
- 4.4. The individual animals should be tested by species specific PCR for Bovine Babesiosis (*B. bovis* and *B. bigemina*), Bovine Anaplasmosis (*A. marginale* and *A. centrale*) and found to be negative immediately prior to shipment.
- 4.5. The animals have to be tested for Paratuberculosis (by CFT / ELISA/delayed type hypersensitivity (DTH) test during thirty (30) days prior to shipment and found to be negative.
- 4.6. Animals have to be tested for Leptospirosis by Microscopic Agglutination Test (MAT) during thirty (30) days prior to shipment and found to be negative.
- 4.7. Animals have to be tested for **Brucellosis** by RBPT/CFT during thirty (30) days prior to shipment and found to be negative.
- 4.8. **Bovine Genital Campylobacteriosis** (*Campylobacter fetus* subsp. *venerealis*) and **Trichomonosis** (*Tritrichomonas foetus*) should fulfil following requirements:  
  
During the pre-export quarantine period the individual animals to be tested by culture of pre-nuptial washings (bulls) or vaginal mucus (heifers/cows) for **Campylobacteriosis** and **Trichomonosis** and found to be negative.
- 4.9. Animals to be tested for **Q fever** with PCR or ELISA (Preferred PCR) 30 days prior to shipment and found to be negative.
- 4.10. The animals should be negative for **Salmonella Dublin** by ELISA when 2 tests performed 2 months apart and the last test performed within 30 days immediately prior to shipment.
- 4.11. The animals to be tested with a recommended sedimentation test to identify liver flukes 14 days after the second treatment, while in pre-export isolation period and should be negative.
- 4.12. The animals show no evidence of Screw worm infestation and skin diseases

- 4.13. Animals should be immunized against Anaplasmosis, Babesia bovis and Babesia bigemina not less than 30 days and not more than 3 months immediately prior to shipment.
- 4.14. The animals to be vaccinated against Leptospirosis (Pomona and hardjo) by using 7 in 1 Leptospirosis vaccine not less than twenty one (21) days and not more than three (3) months prior to embarkation.
- 4.15. The animals to be vaccinated against *Salmonella Dublin* not less than twenty one (21) days and not more than three (3) months prior to embarkation.
- 4.16. The animals to be vaccinated against Clostridial diseases with 5 in 1 vaccine; *C. perfringens*, *C. chauvoei*, *C. septicum*, *C. novyi* and *C. tetani* not less than twenty one (21) days and not more than 3 months prior to embarkation.

## 5. Treatment

- 5.1. The animals to be treated for Cestodes and Nematodes twice with an effective broad spectrum anthelmintic with an interval of 7-10 days during pre-export quarantine period.
- 5.2. The animals to be treated twice at 10-14 days interval for flukes, with a Flukicide effective against all stages of liver Flukes (*Fasciola hepatica*) during the pre-export quarantine period.
- 5.3 The animals to be treated for all ectoparasites within seven (7) days before loading to the Vessel and should be completely free of all ectoparasites.

## 6. Clinical Examination and Transport

- 6.1. The selected animals should be kept separate from other cattle in the herd of origin for at least twenty one (21) days immediately before transferring to a pre-export quarantine station approved by the state Veterinary authority. The cattle should be kept in quarantine station for **minimum of thirty (30) days** immediately prior to the embarkation at the port of origin.
- 6.2. Within 3 (three) days prior to the scheduled date of shipment individual animals should be examined and found to be in good health, with no deformities and fit for transport.
- 6.3. Animals should not show clinical signs of any infectious and contagious diseases,
- 6.4. The animals should be transported under conditions laid down by OIE (chapter 7.2 of Terrestrial Animal Health Code 2019) and International Air /Sea Transport Association (IATA/IASA) regulations.
- 6.5. During the voyage, adequate Veterinary support should be ensured by the accompany of Veterinarians who have been qualified and registered under the veterinary council of India.
- 6.6 All manure, litter and bedding materials in the animal carrying vessels should be disposed of in such a way as to prevent the transmission of diseases and in compliance with all

relevant health and environmental legislation described under OIE specifications and international regulations.

## 7. General Conditions

7.1. Importation of animals must be strictly adhere to the part III ,section 22,23,24,25,26,27 and 28 of Sri Lanka Animal disease act, No.59 of 1992.

7.2. Record of all the test results, vaccinations and treatments done (Name of the vaccine/ drugs and the date of administration) along with all the relevant information to be submitted to the Director General/ Department of Animal Production and Health, Sri Lanka and get the permission 7 days before loading of animals.

7.3. Post import quarantine testing

The following health requirement verification tests will be performed by Veterinary Research Institute of Sri Lanka (The cost of laboratory tests must be borne by the importer).

- i. BVD: RT-PCR (Ag), ELISA (Ab)
- ii. IBR/IPV:by RT-PCR (Ag), ELISA (Ab)
- iii. Blue tongue: by RT-PCR(Ag), ELISA (Ab)
- iv. Brucellosis by RBPT/CFT
- v. Bovine Tuberculosis : by PPD skin test
- vi. Bovine paratuberculosis : CFT
- vii. Campylobacteriosis: by culture, preputial washing & vaginal mucous
- viii. Leptospirosis Microscopic agglutination test & urine culture for the organism.
- ix. Tricomoniasis:- by culture, preputial washing & vaginal mucous
- x. Blood parasites: (Anaplasma, Babesia ) by using PCR
- xi. Examination of dung samples for **Cestodes** and Nematodes eggs – by Faecal flotation concentration test 14 days after second treatment - should be negative immediately prior to embarkation
- xii. Examination of dung samples for **liver flukes** eggs - by Sedimentation concentration test 14days after second treatment - should be negative immediately prior to embarkation

The health requirement verification tests to be done during the 30 days of quarantine period in Sri Lanka and found to be negative before release of imported animals from the post-import quarantine locations. If any animal will fail to fulfil the health requirements stipulated under health requirements or the negative results in testing for above diseases during the in-country quarantine period, animals should be re-exported or destroyed under the provision in Custom ordinance No 17 of 1869 and Animal Disease Act No. 59 of 1992 respectively. (The cost of such re-export or destruction and compensation for the damage should be borne by the importer)

***Veterinary Health Requirements could be varied, from time to time, depending on the animal disease situation of the exporting country.***

Dr. (Mrs.) V.R.N.Munasinghe, Director (VRA)

D.S.S.P.Silva, Director, Livestock Planning and economic division.

Dr.L.W.N. Samaranayake, Director, Animal breeding

Dr.S.Hettiarachchi, Director, Animal health.

Dr. H. Kothalawala. Head, Virology division, VRI.

Dr. (Mrs.) P. S.Fernando. Head, Bacteriology division,, VRI

Dr.(Mrs). P.G.I.D.Amarasiri, Head, Parasitology division, VRI

Dr. Dr.G.G.I.A Jayawickrama ,Veterinary surgeon, Animal Health division

Dr.B.M.M.Ekanayake, Veterinary Surgeon, VRA division